

*Published in Europhysics Lett. 29, 515 (1995)*

## Statistics of Oscillator Strengths in Chaotic Systems

Nobuhiko Taniguchi, A. V. Andreev and Boris L. Altshuler

*Department of Physics, Massachusetts Institute of Technology,*

*77 Massachusetts Avenue, Cambridge, MA 02139*

*and*

*NEC Research Institute, 4 Independence Way, New Jersey 08540*

()

### Abstract

The statistical description of oscillator strengths for systems like hydrogen in a magnetic field is developed by using the supermatrix nonlinear  $\sigma$ -model. The correlator of oscillator strengths is found to have a universal parametric and frequency dependence, and its analytical expression is given. This universal expression applies to quantum chaotic systems with the same generality as Wigner-Dyson statistics.

PACS number: 05.45.+b, 32.60.+i

Typeset using REVTeX

A conventional way to approach the phenomenon of quantum chaos is to study spectra of quantum systems [1]. These spectra are known to be well described by Wigner-Dyson statistics of energy levels [2], based on the eigenvalue distribution of random matrix ensembles. Usually chaos manifests itself in highly excited states of a system. Excitation from a regular state (e.g., the ground state) to a chaotic state can be achieved by applying an external perturbation. Experiments of this type provide *independently* the transition probabilities from a given initial state to excited states, as well as the spectrum of the system. These probabilities for optical excitations are known as oscillator strengths. Like the eigenenergies, the oscillator strengths of highly excited states are random, and should be described statistically. Along this line, the distribution of the transition matrix elements was predicted to have the Porter-Thomas distribution in the chaotic limit [3].

In this Letter we develop a statistical description of oscillator strengths based on the Wigner-Dyson hypothesis. We investigate a statistical characteristic different from Ref. [3] — the correlation function of oscillator strengths at *different* frequencies and “magnetic fields”. It is emphasized that we make no further assumptions such as on the form of the matrix elements. Since the information contained within the oscillator strengths is independent of the spectral properties, their statistics provide a characterization of quantum chaos which complements the usual energy level correlations. We determine universal parametric correlations of oscillator strengths which should apply with the same generality as Wigner-Dyson statistics. We have also discovered an unexpected differential relation between this parametric correlations and the parametric density of states correlator of Ref. [4].

The results of this Letter can be applied to responses of various impurity states in semiconductors, or to microwave and optical excitations of chaotic quantum dots etc. They can be easily modified to describe a much broader class of phenomena, e.g., cross sections of nuclear excitations by neutrons. However, as a concrete example, we have in mind the hydrogen atom in a magnetic field, which attracts great interest as a

“real” quantum system with a chaotic classical analogue. The energy spectrum of this system is known to obey Wigner-Dyson statistics [5]. In addition, the distribution of the transition matrix elements is observed to obey the Porter-Thomas distribution [6].

The oscillator strength per unit energy is conventionally defined as

$$W(\bar{\Omega}, \bar{B}) = 2\bar{\Omega} \sum_{\alpha} |\langle \Phi_0 | \hat{z} | \alpha \rangle|^2 \delta(E_{\alpha}(\bar{B}) - E_0(\bar{B}) - \bar{\Omega}), \quad (1)$$

where  $B$  denotes the magnetic field. More generally,  $B$  can stand for any external parameter of the system. In writing down Eq. (1), we use the transition matrix element between a certain initial state  $|\Phi_0\rangle$  with the energy  $E_0$  and a highly excited state  $|\alpha\rangle$  with  $E_{\alpha}$  induced by the light polarized along  $z$ -axis. However, it is not necessary to assume this particular form of the transition matrix elements to proceed our theory.

We have evaluated the correlation function of oscillator strengths

$$S(\Omega, B) = \langle W(\bar{\Omega}, \bar{B}) W(\bar{\Omega} + \Omega, \bar{B} + B) \rangle, \quad (2)$$

for small energy separation ( $\Omega \ll \bar{\Omega}$ ) and at slightly different “magnetic fields” ( $B \ll \bar{B}$ ). Here  $\langle \dots \rangle$  denotes the averaging over a certain interval of the spectrum and/or of the parameter  $\bar{B}$ . The magnitude of the correlator  $S(\Omega, B)$  is not universal. It depends on the initial state  $|\Phi_0\rangle$ , on the type of excitation, and on the direction of the polarization. (See Eq. (17) below.) However, the dependence of  $S$  on  $\Omega$  and  $B$  turns out to be determined by a universal function  $s(\omega, x)$ ,

$$S(\Omega, B) = S_0 [s(\omega, x) + 1]; \quad S_0 = \langle W \rangle^2, \quad (3)$$

where we use the rescaled parameters defined by [4]

$$\omega \equiv \Omega / \Delta; \quad \Delta = \langle E_{\alpha+1} - E_{\alpha} \rangle, \quad (4)$$

$$x \equiv B \sqrt{C(0)}; \quad C(0) = \langle (\partial E_{\alpha} / \partial B)^2 \rangle / \Delta^2. \quad (5)$$

The two terms in Eq. (3) correspond to the connected and disconnected parts of the correlator. The function  $s(\omega, x)$  is classified according to the Dyson universality

classes: orthogonal ( $o$ ), unitary ( $u$ ), or symplectic ( $s$ ). It is convenient to present  $s(\omega, x)$  by

$$s_e(\omega, x) = \left( \frac{\partial^2}{\partial \omega^2} + 2\zeta_e \frac{\partial}{\partial x^2} \right) h_e(\omega, x), \quad (6)$$

where  $e = o, u, s$ ;  $\zeta_o = \zeta_s = 2$ ,  $\zeta_u = 1$  and

$$h_u(\omega, x) = \frac{-1}{2\pi^2} \int_{-1}^1 d\lambda \int_1^\infty d\lambda_1 \frac{\cos[\pi\omega(\lambda_1 - \lambda)]}{(\lambda_1 - \lambda)^2} \exp\left[-\frac{\pi^2 x^2}{2}(\lambda_1^2 - \lambda^2)\right], \quad (7)$$

$$h_{o,s}(\omega, x) = \frac{-1}{\pi^2} \iiint d\lambda d\lambda_1 d\lambda_2 \frac{|1 - \lambda^2| \cos[\pi\omega(\lambda_1 \lambda_2 - \lambda)]}{(\lambda_1^2 + \lambda_2^2 + \lambda^2 - 2\lambda_1 \lambda_2 \lambda - 1)^2} \times \exp\left[-\frac{\pi^2 x^2}{4}|2\lambda_1^2 \lambda_2^2 - \lambda_1^2 - \lambda_2^2 - \lambda^2 + 1|\right]. \quad (8)$$

In Eq. (8), the integral region is defined by  $\lambda \in (-1, 1)$ ,  $\lambda_{1,2} \in (1, \infty)$  for the orthogonal case, or  $\lambda \in (1, \infty)$ ,  $\lambda_1 \in (-1, 1)$ , and  $\lambda_2 \in (0, 1)$  for the symplectic case.

Let us compare  $S(\Omega, B)$  with the more conventional statistical characteristic of quantum chaotic systems. For a certain value of  $B$ , the density of states at energy  $E$  can be written as  $\nu(E, B) = \sum_\alpha \delta(E - E_\alpha(B))$ . The parametric correlator of densities of states is written in terms of rescaled variables Eqs. (4,5)

$$\langle \nu(E, \bar{B}) \nu(E + \Omega, \bar{B} + B) \rangle = \frac{1}{\Delta^2} [1 + k(\omega, x)] \quad (9)$$

The explicit forms for  $k_o(\omega, 0)$ ,  $k_u(\omega, 0)$  and  $k_s(\omega, 0)$  were obtained by Dyson (see, e.g., Ref. [2]), and were recently evaluated at finite  $x$  [4] by the supermatrix method [7]. In our notation these results can be written as [8]

$$k(\omega, x) = \partial^2 h(\omega, x) / \partial \omega^2. \quad (10)$$

Both  $k(\omega, x)$  and  $s(\omega, x)$  are singular at  $\omega = 0$  and  $x = 0$ . We present in Fig. 1 the ratio  $r(\omega, x) = (1 + s(\omega, x)) / (1 + k(\omega, x))$  which is not singular.

Comparing  $s(\omega, x)$  of Eq. (6) with  $k(\omega, x)$  of Eq. (10), we get a differential relation:

$$\frac{\partial^2}{\partial \omega^2} s(\omega, x) = \left( \frac{\partial^2}{\partial \omega^2} + 2\zeta \frac{\partial}{\partial x^2} \right) k(\omega, x). \quad (11)$$

This relation is not trivial at all [8], especially when considered from the semiclassical point of view [1,9]. That approach is based on the approximation of the density of states  $\nu(E, B)$  by a certain sum over the classical periodic trajectories. Although the averaged oscillator strength itself may be expressed by the summation of the periodic trajectories [10], the oscillator strength correlator apparently requires all *closed* trajectories to be taken into account, not just the classical *periodic* ones. (See also Ref. [11] for some discussions.) Eqs. (6,10) for  $s(\omega, x)$  and  $k(\omega, x)$  are valid for arbitrary  $\omega$  and  $x$ , including nonperturbative regime at  $\omega \lesssim 1$ . We believe that these formulas as well as Eq. (11) have wider region of applicability than the conventional semiclassical approach.

It is worth mentioning that the definition of  $h(\omega, x)$  should be interpreted rather formally, and in fact, integral expressions Eq. (7,8) for  $h(\omega, x)$  diverge logarithmically. Of course, the corresponding integrals for  $s(\omega, x)$  and  $k(\omega, x)$  are converged. However, we have introduced  $h(\omega, x)$  not only to make formulas for  $s(\omega, x)$  and  $k(\omega, x)$  more compact. Note that the finite part of  $h(\omega, x) - h(0, x)$  is related to the generalized parametric number variance function  $v(\omega, x)$  by (compare with Ref. [12])

$$h(\omega, x) - h(0, x) = \frac{1}{2} (v(\omega, x) - v(0, x)), \quad (12)$$

where  $v(\omega, x) = [n(\epsilon + \omega, x + \bar{x}) - n(\epsilon, \bar{x}) - \omega]^2$  and  $n(\epsilon, x) = \sum_i \theta(\epsilon - \epsilon_i(x))$ .

Now let us sketch the derivation of our main results (3-8). One can write  $S(\Omega, B)$ ,

$$S(\Omega, B) = \int d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 A(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4) R(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4), \quad (13)$$

where

$$A = \frac{4\bar{\Omega}^2}{\pi^2} z_1 z_2 z_3 z_4 \Phi_0^*(\mathbf{r}_1) \Phi_0(\mathbf{r}_2) \Phi_0^*(\mathbf{r}_3) \Phi_0(\mathbf{r}_4), \quad (14)$$

$$R = \left\langle \text{Im} G_{E_0 + \bar{\Omega}}^R(\mathbf{r}_1, \mathbf{r}_2; \bar{B}) \text{Im} G_{E_0 + \bar{\Omega} + \Omega}^R(\mathbf{r}_3, \mathbf{r}_4; \bar{B} + B) \right\rangle. \quad (15)$$

and  $G_E^{R,A}(\mathbf{r}, \mathbf{r}'; B) = \langle \mathbf{r} | (E - H(B) \pm i0)^{-1} | \mathbf{r}' \rangle$  are the retarded and advanced Green's functions. The factor  $A$  is determined by the initial state wave function  $\Phi_0(\mathbf{r})$ , whereas

$R$  reflects only the properties of excited states, which are presumed chaotic. In contrast to the evaluation of  $K(\Omega, B)$  in Eq. (9), we now have to accommodate the dependence on four different positions  $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4$  as depicted in Fig. 2a.

We apply the zero-dimensional supermatrix model [7] to determine  $R$ :

$$\begin{aligned} \frac{2}{\pi^2 \nu^2} R(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4) = & f(r_{12})f(r_{34}) \text{Re} \langle 1 - Q_{33}^{11} Q_{33}^{22} \rangle_Q \\ & - f(r_{13})f(r_{24}) \text{Re} \langle Q_{34}^{12} Q_{43}^{21} \rangle_Q - f(r_{14})f(r_{32}) \text{Re} \langle Q_{33}^{12} Q_{33}^{21} \rangle_Q, \end{aligned} \quad (16)$$

where the dimensionless function  $f(r) = \langle \text{Im} G^R(r) \rangle / \langle \text{Im} G^R(0) \rangle$  which is not universal [14]. For example,  $f(r)$  becomes  $\sin(pr) \exp(-r/2\ell)/pr$  in three dimensional metallic grain with the Fermi momentum  $p$  and the mean free path  $\ell$ .  $\langle \dots \rangle_Q$  of Eq. (16) denotes the integration over the saddle point manifold  $\langle \dots \rangle_Q \equiv \int DQ (\dots) \exp(-(\pi\omega/4) \text{STr} Q \Lambda - (\pi^2 x^2/32\zeta_e) \text{STr}[Q, \Lambda]^2)$ , and the definition of  $\text{STr}$  as well as  $\Lambda$  and the structure of supermatrices  $Q$  can be found in [7]. We can identify physically each term of Eq. (16) as in Fig. 2(b-d), — DOS fluctuation-type (Fig. 2b), Cooperon-type (Fig. 2c) and Diffuson-type (Fig. 2d). In fact, we get the DOS correlator of Eq. (9) by integrating Fig. 2b over  $\mathbf{r}_1$  and  $\mathbf{r}_3$ , whereas integrating Fig. 2c and 2d over  $\mathbf{r}_1$  and  $\mathbf{r}_3$  (Fig. 2c) or  $\mathbf{r}_1, \mathbf{r}_2$  (Fig. 2d) gives another function which was recently shown to correspond to the dielectric function of periodic systems with chaotic unit cell [13]. Definite integration over  $Q$ -matrix leads to  $\text{Re} \langle Q_{33}^{11} Q_{33}^{22} - 1 \rangle_Q = -2 - 2\partial^2 h(\omega, x)/\partial\omega^2$ ,  $\text{Re} \langle Q_{33}^{12} Q_{33}^{21} \rangle_Q = -4 \partial h(\omega, x)/\partial x^2$ , and the Cooperon-type contribution  $\langle Q_{34}^{12} Q_{43}^{21} \rangle_Q$  is equal to the Diffuson-type  $\langle Q_{33}^{12} Q_{33}^{21} \rangle_Q$  for the orthogonal and symplectic cases, or vanishes for the unitary. Everything combined, we obtain our results (3-8), with the prefactor  $S_0$  equal to

$$S_0 = \left[ 2\bar{\Omega}\nu \int d\mathbf{r}_1 d\mathbf{r}_2 z_1 z_2 \Phi_0^*(\mathbf{r}_1) f(r_{12}) \Phi_0(\mathbf{r}_2) \right]^2, \quad (17)$$

since  $\Phi_0(\mathbf{r})$  is real for orthogonal and symplectic cases. As it was mentioned, the constant  $S_0$  as well as the parameters  $\Delta$  and  $C(0)$  is not universal. However, the dependence of  $S$  on the rescaled frequency  $\omega$  and “magnetic field”  $x$  is universal.

In conclusion, we have developed the statistical description of oscillator strengths of a chaotic system such as hydrogen in a magnetic field. In particular, we have obtained the analytical form of the correlator of oscillator strengths  $S(\Omega, B)$ , dependent on  $\Omega$  and  $B$  (Eqs. (3-6)). We have found an identity Eq. (11) that relates  $S$  with the density of states correlator. It looks like a challenge for the semiclassical theory of quantum chaos to derive this differential relation. The universality of  $S(\Omega, B)$  is as general as that of the correlation function of densities of states  $K(\Omega, B)$ , and should be applicable to various physical situations: impurity states in semiconductors, to microwave and optical excitations in chaotic quantum dots, etc. We believe that experimental studies of the statistics of the oscillator strength provides a new way of characterizing the chaotic systems, complimentary to the spectral statistics.

The authors are especially grateful to B. D. Simons for valuable discussions. We also like to thank M. Courtney, D. Kleppner, E. Mucciolo, and V. N. Prigodin for helpful discussions. The work was partially supported by Joint Services Electronic Program Contract No. DAAL 03-89-0001. N.T. also acknowledges partial support from the research fellowship from Murata Overseas Scholarship Foundation.

## REFERENCES

- [1] M. Berry, in *Chaos and Quantum Physics*, edited by M.-J. Giannoni, A. Voros, and J. Zinn-Justin (North-Holland, Amsterdam, 1991), pp. 251.
- [2] M. L. Mehta, *Random Matrices 2nd ed.* (Academic Press, San Diego, 1991).
- [3] Y. Alhassid, and R. D. Levine, Phys. Rev. Lett. **57**, 2879 (1986).
- [4] B. D. Simons and B. L. Altshuler, Phys. Rev. B **48**, 5422 (1993).
- [5] D. Wintgen and H. Friedrich, Phys. Rev. Lett. **57**, 571 (1986); D. Delande and J. C. Gray, *ibid.*, 2006; G. Wunner *et. al.*, *ibid.*, 3261. For a review with recent results, see D. Delande, in *Fundamental Systems in Quantum Optics*, edited by J. Dalibard, J. M. Raimond, and J. Zinn-Justin (North-Holland, Amsterdam, 1992), pp. 381; D. Kleppner, *ibid.*, pp. 417.
- [6] G. Wunner, W. Schweizer, and H. Ruder, in *The Hydrogen Atom* (Springer-Verlag, Berlin, 1989), pp. 300.
- [7] K. B. Efetov, Adv. Phys. **32**, 53 (1983). See also J. J. M. Verbaarschot, H. A. Weidenmüller, and M. R. Zirnbauer, Phys. Rep. **129**, 367 (1985).
- [8] As was shown by B. D. Simons *et.al.*, Nucl. Phys. **B409**, 487 (1993),  $k(\omega, x) = \partial^2 h / \partial \omega^2$  gives the density-density correlator for the one-dimensional gas with  $1/r^2$  interaction when  $\omega \rightarrow r$  and  $-ix^2/2 \rightarrow t$ . In the same sense, after the Fourier transformation in  $r$  and  $t$ ,  $\partial h / \partial x^2$  is found to give the dynamical conductivity  $\sigma(q, \omega)$  of this one-dimensional gas with the help of the continuity relation.
- [9] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer-Verlag, New York, 1990).
- [10] M. Wilkinson, J. Phys. A **20**, 2415 (1987).
- [11] E. B. Bogomol'nyi, Sov. Phys. JETP **69**, 275 (1989).



- [12] J. Goldberg *et al.*, Nonlinearity **4**, 1 (1991); Goldberg and Schweizer, J. Phys. A **24**, 2785 (1991).
- [13] N. Taniguchi and B. L. Altshuler, Phys. Rev. Lett. **71**, 4031 (1993).
- [14] A similar formula was used in V. N. Prigodin *et.al.*, Phys. Rev. Lett. **72**, 546 (1994) to study the time evolution of the density of quantum dots .

# FIGURES




FIG. 1. Schematic plot of  $1 + s(\omega, x)$  normalize by  $1 + k(\omega, x)$  for orthogonal (above) and unitary (below) cases.




FIG. 2. (a) Diagram representation of  $R(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4)$ . (b) DOS fluctuation-type contribution. (c) Cooperon-type contribution. (d) Diffuson-type contribution.